SHP-77 Cell Line

Cat. No. : CL-0718

Origin and General Characteristics

Cell Name	SHP-77
Organism	Human
Age	54Y
Sex	Male
Tissue	Lung
Morphology	Round
Growth Properties	Suspension, clustered; loosely adherent
Background	SHP-77 retains important characteristics of small cell lung cancer (SCLC) and possesses stable biochemical properties, allowing continuous cultivation. This cell line is an unusual undifferentiated large-cell variant of SCLC. Although it exhibits a variant morphology, it maintains the biochemical properties of classical SCLC. Electron microscopy reveals gland formation and the presence of intracytoplasmic lamellar bodies. These cells express the neuroendocrine marker L-dopa decarboxylase and contain dense-core secretory granules. SHP-77 can serve as a target for detecting cytotoxicity via ^{51}Cr and ^{111}In release in both in vitro assays and in vivo nude mouse models, and is used to evaluate the immune reactivity of cells and serum from lung cancer patients. This cell line can also be used to assess the immune status of SCLC patients undergoing radiotherapy or chemotherapy.
Biosafety Level	BSL-1

Culture Conditions and Handling

Complete Medium	RPMI-1640[PM150110]+10% Nutrient+1% Supplement
Subcultivation Ratio	4-5×10^5 cells/mL
Medium Renewal	2 to 3 times per week
Incubation Atmosphere	Air, 95%; CO_2, 5%
Temperature	37°C
Freeze Medium	General Freezing Medium[PB180436]
Storage Conditions	For long-term cryopreservation, cryovials should be stored in liquid nitrogen at −150°C to −196°C. Storage at −80°C is restricted to short-term interim use only.
Instructions	1. Check all containers for leakage or breakage. 2. Remove the frozen cells from the dry ice packaging and immediately transfer them to liquid nitrogen (liquid or vapor phase) for long-term cryopreservation.

Subculturing Procedure	1. This cell line exhibits mixed adherent and suspension growth phenotypes. Suspended cells remain viable and must be retained. Transfer the cell suspension to a centrifuge tube and centrifuge at 1200 rpm (approximately 250 ×g). 2. Adherent cells can be detached by gentle pipetting to obtain a homogeneous cell suspension. 3. Combine all harvested suspension and adherent cell populations, resuspend in fresh complete growth medium, and seed into new culture vessels at the recommended subcultivation ratio. Loosen the cap or use a vented cap for incubation.

Special Features of the Cell Line

Tumorigenic	Yes; Yes, the cells form tumors in athymic nude mice, and usually grow as circumscribed nodules without evidence of metastases.
Antigen Expression	Blood Type O; Rh +; CD56; CD57 (HNK-1,Leu-7)
Genes Expressed	neural cell adhesion molecule (NCAM) NKH-1

Handling Recommendations for Cryopreserved Cells

Preparations Before Thawing

1. Prepare the recommended complete culture medium and preheat it to 37°C for 30 minutes.
2. Prepare 9 mL of complete medium within a sterile centrifuge tube to dilute the cryoprotectant present in the frozen cell suspension.
3. Use sterile gloves to protect the cryovial from contact with the water bath water, and preventing cell contamination.

Thawing Procedure

1. Retrieve the cryovial containing frozen cells from liquid nitrogen storage. Place it in sterile gloves and immediately submerge it in a 37°C water bath.
2. Thaw the cells rapidly (<1 minute) by gently swirling the vial in the 37°C water bath until only a small ice crystal remains.
3. Transfer the vial to a biosafety cabinet. Wipe the exterior of the vial with 75% ethanol before opening.
4. Transfer the thawed cells dropwise into a preheated 9 mL centrifuge tube containing complete medium.
5. Centrifuge the cell suspension at approximately 250 × g for 5 minutes.
6. After centrifugation, discard the supernatant and retain the cell pellet.
7. Gently resuspend the cells in complete growth medium, then transfer them to an appropriate culture vessel and place it in the recommended culture environment.

Notes

1. The entire recovery process should be completed as quickly as possible.
2. Select the culture vessel size based on the number of cells in the cryovial. Procell single cryovials are recommended for resuspension in T25 flasks or 10 cm dishes.
3. Minimize the time thawed cells are kept at room temperature. Cryoprotectant must be immediately diluted or removed by centrifugation.

Precautions After Receipt of Frozen Cells

1. Upon unpacking, inspect the condition of the frozen cells and dry ice, and take photographs immediately. The following after-sales service will be provided based on these photographs, including assessment of the remaining dry ice, verification that the cryovial was fully buried in dry ice, and evaluation of whether the cells thawed and refrozen during transit.
2. Upon receipt, the cells should be transferred to liquid nitrogen immediately or directly resuscitated. If liquid nitrogen is unavailable, the cells can be temporarily stored at -80°C, however, the storage period should be limited to less than two weeks whenever possible. Prolonged storage at -80°C may gradually reduce post-thaw cell viability, and the extent of this viability loss is unpredictable.
3. Ensure that the operator has sufficient knowledge and experience in cell culture, and the laboratory is equipped with essential instruments, including a biosafety cabinet, CO_2 incubator, inverted microscope, centrifuge, water bath. Carefully review the cell instruction manual to understand key cell characteristics including growth properties (adherent or suspension), morphology, basal medium requirements, serum concentration, cytokine supplementation, subcultivation ratio, medium renewal schedule.
4. After resuscitation, observe the cells under a microscope and record the cell status by taking photographs (1-3 images each at 100× and 200× magnification for 3 consecutive days). These images will serve as supporting documentation for follow-up services. In addition, a small aliquot of cells may be used to assess cell viability by automated cell counting or trypan blue staining.

Note: Cells should not be observed too frequently within the first 24 hours after resuscitation, as this may affect cell growth or adherence. Observation once per day is sufficient.

5. After sufficient expansion, the cells should be cryopreserved as soon as possible to ensure availability of backup stocks for future experiments or in case of unexpected experimental failure. Cells in the logarithmic growth phase should be selected for cryopreservation. After cryopreservation, it is recommended to resuscitate one vial to verify post-thaw viability and confirm successful cryopreservation.

Procell system®
Procell system®
Procell system®