Elabscience®

Recombinant Rabbit B7-2/CD86 Protein (His Tag)

Catalog Number: PKSQ050067

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Rabbit		
Source	HEK293 Cells-derived Rabbit Rabbit B7-2/CD86 protein Ala23-Arg246, with an C-		
	terminal His		
Calculated MW	26.3 kDa		
Observed MW	38-70 kDa		
Accession	P42071		
Bio-activity	Not validated for activity		
Properties			
Purity	>95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

kDa 120 90	MK	R
60		
40	-	100
30	-	
20	-	
14	-	

> 95 % as determined by reducing SDS-PAGE.

Background

T-lymphocyte activation antigen CD86 (B7-2) is a glycosylated protein in the B7 family. B7 family members are transmembrane cell surface molecules that play important roles in immune activation and the maintenance of immune tolerance. It is highly expressed on activated antigen presenting cells. CD86 involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. It may play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. It is expressed by activated B-lymphocytes and monocytes and promoted by MARCH8 and results in endocytosis and lysosomal degradation.

For Research Use Only

Tel:400-999-2100