

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® Violet 450 Anti-Human CD15/SSEA-1 Antibody[HI98]

Catalog Number: E-AB-F1079Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human Mouse Host Isotype Mouse IgM, ĸ HI98

Clone No.

Isotype Control Elab Fluor® Violet 450 Mouse IgM, κ Isotype Control[MM-30] [Product E-AB-F09782Q]

Conjugation Elab Fluor® Violet 450

**Conjugation Information** Elab Fluor® Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

#### **Applications** Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Human peripheral blood granulocytes are stained with Elab

Fluor® Violet 450 Anti-Human CD15 Antibody (filled gray histogram) or Elab Fluor® Violet 450 Mouse IgM, κ Isotype Control (empty black histogram).

#### **Preparation & Storage**

Keep as concentrated solution. **Storage** 

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

#### **Antigen Information**

**Alternate Names** 3-FAL;3-FL;LNFP III;Lewis X;LexSSEA-1;X-hapten;SSEA-1

**Uniprot ID** P22083 Gene ID 2526

# For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Background**

CD15 is 3-fucosyl-N-acetyllactosamine (3-FAL), also known as Lewis X, 3-FAL, Xhapten, and SSEA-1. CD15 is expressed on granulocytes and monocytes. It has also been shown to be expressed on Langerhans cells and some malignant cells. CD15 has been implicated in adhesion, as well as chemotaxis, phagocytosis, and bactericidal activity.

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623  $Web: \underline{w\ w\ w\ .elabscience.com}$