

## NUMA/NMP22 Monoclonal Antibody(Capture)

catalog number: **AN001180P**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human NUMA/NMP22 protein expressed by E.coli
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG2a
<b>Clone</b>	2F1
<b>Purification</b>	Protein A/G Purification
<b>Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.05% Proclin300.

### Applications Recommended Dilution

<b>ELISA Capture</b>	2-8 µg/mL
----------------------	-----------

### Data



Sandwich ELISA-Recombinant Human NUMA/NMP22 protein standard curve. Background subtracted standard curve using NUMA/NMP22 antibody(AN001180P) (Capture), NUMA/NMP22 antibody(AN001190P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human NUMA/NMP22 protein is 0.625-40 ng/mL.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

### Background

### For Research Use Only

Microtubule (MT)-binding protein that plays a role in the formation and maintenance of the spindle poles and the alignment and the segregation of chromosomes during mitotic cell division. Functions to tether the minus ends of MTs at the spindle poles, which is critical for the establishment and maintenance of the spindle poles. Plays a role in the establishment of the mitotic spindle orientation during metaphase and elongation during anaphase in a dynein-dynactin-dependent manner. In metaphase, part of a ternary complex composed of GPSM2 and G(i) alpha proteins, that regulates the recruitment and anchorage of the dynein-dynactin complex in the mitotic cell cortex regions situated above the two spindle poles, and hence regulates the correct orientation of the mitotic spindle. During anaphase, mediates the recruitment and accumulation of the dynein-dynactin complex at the cell membrane of the polar cortical region through direct association with phosphatidylinositol 4,5-bisphosphate (PI(4,5)P2), and hence participates in the regulation of the spindle elongation and chromosome segregation. Binds also to other polyanionic phosphoinositides, such as phosphatidylinositol 3-phosphate (PIP), lysophosphatidic acid (IPA) and phosphatidylinositol triphosphate (PIP3), in vitro. Also required for proper orientation of the mitotic spindle during asymmetric cell divisions. Plays a role in mitotic MT aster assembly. Involved in astral spindle assembly. Positively regulates TNKS protein localization to spindle poles in mitosis. Highly abundant component of the nuclear matrix where it may serve a non-mitotic structural role, occupies the majority of the nuclear volume. Required for epidermal differentiation and hair follicle morphogenesis.