

Recombinant Mouse CLEC3B/Tetranectin Protein (His Tag)

Catalog Number: PKSM040537

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse CLEC3B/Tetranectin protein Met 1-Val 202, with an C-terminal His
Calculated MW	20.7 kDa
Observed MW	25 kDa
Accession	P43025
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Tetranectin (TN), also known as C-type lectin domain family 3, member B (CLEC3B) is a member of the C-type lectin Family. It is plasminogen kringle 4 binding protein and regulates fibrinolysis and proteolytic processes via binding to plasminogen. Tetranectin has been suggested to play a role in tissue remodeling, due to its ability to stimulate plasminogen activation and its expression in developing tissues such as developing bone and muscle. Tetranectin enhances plasminogen activation by a tissue-type plasminogen activator so that it has been suggested to play a role in tissue remodeling. Tetranectin may play a role in the wound healing process. Tetranectin may play a role in neurological diseases and may serve as a diagnostic aid in multiple sclerosis (MS). Tetranectin was found significantly under-expressed in both serum and saliva of metastatic oral squamous cell carcinoma (OSCC) compared to primary OSCC. Tetranectin is thought to enhance proteolytic processes enabling tumor cells to invade and metastasize.