

A Reliable Research Partner in Life Science and Medicine

PerCP Anti-Human CD4 Antibody[RPA-T4]

Catalog Number: E-AB-F1109F

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ **Clone No.** RPA-T4

Isotype Control PerCP Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792F]

Conjugation PerCP

Conjugation Information PerCP is designed to be excited by the blue laser (488 nm) and detected using an optical

filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with PerCP Anti-Human CD4 Antibody[RPA-T4] (filled gray histogram) or PerCP Mouse lgG1, κ Isotype Control (empty black histogram). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4

Web: www.elabscience.cn

 Uniprot ID
 P01730

 Gene ID
 920

For Research Use Only

Elabscience®

Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD4, also known as T4/Leu-3, is a 55 kD single-chain type I transmembrane glycoprotein and member of the immunoglobulin superfamily. It is expressed on most thymocytes, helper T cells, type II NKT cells, and monocytes/macrophages. CD4 is part of the TCR/CD3 complex, binds to $\beta2$ domain from the MHC class II molecule, and participates in TCR signal transduction. CD4 is the receptor of IL-16 and is a coreceptor for the human immunodeficiency virus (HIV) and human herpes virus 7 (HHV-7).