

A Reliable Research Partner in Life Science and Medicine

## Recombinant E-Cadherin/CDH1/E-cad/CD324 Monoclonal Antibody

catalog number: AN300024P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human

Immunogen Recombinant Human E-Cadherin / CDH1 / E-cad / CD324 Protein


HostRabbitIsotypeIgGCloneA1134PurificationProtein A

Buffer 0.2 µm filtered solution in PBS

#### Applications Recommended Dilution

WB 1:500-1:2000IP 4-8 μL/mg of lysate

#### Data



Immunoprecipitation analysis using 4 µL anti-E-cad Monoclonal Antibody and 15 µl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using E-cad Monoclonal Antibody at a dilution of 1:100. Lane A:0.5 mg MCF-7 Whole Cell Lysate, Lane B:0.5 mg A431

Whole Cell Lysate
Observed-MW:130 kDa
Calculated-MW:97 kDa



Western Blot with E-Cadherin / CDH1 / E-cad / CD324 Monoclonal Antibody at dilution of 1:500. Lane A: MCF7 Whole Cell Lysate, Lane B: A431 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

> Observed-MW:130 kDa Calculated-MW:97 kDa

#### **Preparation & Storage**

**Storage** This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

inpping icc bi

**Background** 

### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Epithelial (E) - Cadherin (ECAD), also known as cell-CAM120/80 in the human, uvomorulin in the mouse, Arc-1 in the dog, and L-CAM in the chicken, is a member of the cadherin family of cell adhesion molecules. Cadherins are calcium-dependent transmembrane proteins, which bind to one another in a homophilic manner. On their cytoplasmic side, they associate with the three catenins, alpha, beta, and gamma (plakoglobin). This association links the cadherin protein to the cytoskeleton. Without association with the catenins, the cadherins are non-adhesive. Cadherins play a role in development, specifically in tissue formation. They may also help to maintain tissue architecture in the adult. E-Cadherin may also play a role in tumor development, as loss of E-Cadherin has been associated with tumor invasiveness. E-Cadherin is a classical cadherin molecule. Classical cadherins consist of a large extracellular domain which contains DXD and DXNDN repeats responsible for mediating calcium-dependent adhesion, a single-pass transmembrane domain, and a short carboxy-terminal cytoplasmic domain responsible for interacting with the catenins. E-Cadherin contains five extracellular calcium-binding domains of approximately 110 amino acids each.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2