

Recombinant PAX8 Monoclonal Antibody

catalog number: AN301620L

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant human PAX8 fragment
Host	Rabbit
Isotype	Rabbit IgG
Clone	C141243
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications


Recommended Dilution

IHC	1:50-1:100
IF	1:50


Data




Immunohistochemistry of paraffin-embedded Human thyroid using PAX8 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Mouse thyroid using PAX8 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Rat thyroid using PAX8 Monoclonal Antibody at dilution of 1:100.



Immunofluorescent analysis of (4% Paraformaldehyde) fixed SK-OV-3 cells using anti-PAX8 Monoclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

PAX8 is one of nine transcription factors within the PAX gene family. It has been observed to be expressed during embryonic development however expression attenuates after development in all but a few tissues. Research has indicated that over-expression of PAX proteins in general are not an initiating event in tumorigenesis however they may facilitate malignant development. PAX8 is expressed in 80-96% of all High-grade serous carcinomas (HGSC) and is commonly used as a marker for the disease. In addition PAX8 has become a useful biomarker for differentiating between mullerian mucinous and non-mullerian tumors. PAX8 has also been associated with neoplasms in the kidney and thyroid.