Elabscience®

CREBBP Polyclonal Antibody

catalog number: E-AB-19276

Note: Centrifuge before opening to ensure complete recovery of vial contents.

DescriptionReactivityHuman;Mouse;RatImmunogenSynthetic peptide of human CREBBPHostRabbitIsotypeIgGPurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% ght	
ImmunogenSynthetic peptide of human CREBBPHostRabbitIsotypeIgGPurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% ght	
HostRabbitIsotypeIgGPurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% gh	
IsotypeIgGPurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% gl	
PurificationAntigen affinity purificationConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% gly	
ConjugationUnconjugatedBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% gl	
Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% gb	
	ycerol.
Applications Recommended Dilution	
IHC 1:50-1:200	
Data	
Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using CREBBP Polyclonal Antibody at dilution of 1:60(×200)	-
Preparation & Storage	
Store at -20°C Valid for 12 months, Avoid freeze / thaw cycles.	
StorageStore at -20°C Valid for 12 months. Avoid freeze / thaw cycles.ShippingThe product is shipped with ice pack,upon receipt,store it immediately at the	

Background

This gene is ubiquitously expressed and is involved in the transcriptional coactivation of many different transcription factors. First isolated as a nuclear protein that binds to cAMP-response element binding protein (CREB), this gene is now known to play critical roles in embryonic development, growth control, and homeostasis by coupling chromatin remodeling to transcription factor recognition. The protein encoded by this gene has intrinsic histone acetyltransferase activity and also acts as a scaffold to stabilize additional protein interactions with the transcription complex. This protein acetylates both histone and non-histone proteins. This protein shares regions of very high sequence similarity with protein p300 in its bromodomain, cysteine-histidine-rich regions, and histone acetyltransferase domain. Mutations in this gene cause Rubinstein-Taybi syndrome (RTS). Chromosomal translocations involving this gene have been associated with acute myeloid leukemia. Alternative splicing results in multiple transcript variants encoding different isoforms.