



MEF2D Polyclonal Antibody

catalog number: E-AB-53508

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Synthetic peptide of human MEF2D
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:35-1:200

Data	
	
Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using MEF2D Polyclonal Antibody at dilution of 1:50(×200)	Immunohistochemistry of paraffin-embedded Human tonsil tissue using MEF2D Polyclonal Antibody at dilution of 1:50(×200)

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background	
<p>This gene is a member of the myocyte-specific enhancer factor 2 (MEF2) family of transcription factors. Members of this family are involved in control of muscle and neuronal cell differentiation and development, and are regulated by class II histone deacetylases. Fusions of the encoded protein with Deleted in Azoospermia-Associated Protein 1 (DAZAP1) due to a translocation have been found in an acute lymphoblastic leukemia cell line, suggesting a role in leukemogenesis. The encoded protein may also be involved in Parkinson disease and myotonic dystrophy. Alternative splicing results in multiple transcript variants.</p> <p>MEF2D (Myocyte Enhancer Factor 2D) is a Protein Coding gene. Among its related pathways are Phospholipase-C Pathway and Development Angiotensin activation of ERK. GO annotations related to this gene include transcription factor activity, sequence-specific DNA binding and RNA polymerase II transcription factor activity, sequence-specific DNA binding. An important paralog of this gene is MEF2A.</p>	

For Research Use Only