A Reliable Research Partner in Life Science and Medicine

# Recombinant Human c-KIT/CD117 Protein (His Tag)

Catalog Number: PKSH033592

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Species Human

**Source** HEK293 Cells-derived Human c-KIT/CD117 protein Gln26-Thr520, with an C-terminal

His

Calculated MW56.5 kDaObserved MW150 kDaAccessionP10721

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

#### **Background**

C-Kit/SCF R is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). c-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains and 1 protein kinase domain. It belongs to the protein kinase superfamily and CSF-1/PDGF receptor subfamily. SCF R expression on mast cells enables them to infiltrate SCF-secreting tumors where they promote tumor growth and induce local immune suppression. SCF R is up-regulated on dendritic cells by Th2-orTh17-biasing stimuli, and it is required for subsequent dendritic cell induction of Th2 and Th17 response s. SCF R protects vascular smooth muscle cells from apoptosis and assists in the recovery of cardiac function following myocardial infarction.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017