Recombinant Mouse II17F Protein(Sumo Tag)

Catalog Number: PDEM100141



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 10	00	cri	m	17	٦m
u		\mathbf{u}		7 U	

Species Mouse

Source E.coli-derived Mouse II17f protein Arg29-Ala161, with an N-terminal Sumo

 Mol_Mass
 27.5 kDa

 Accession
 Q7TNI7

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse II17f proteins, 2 µg/lane of Recombinant Mouse II17f proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 27.5 KD

Background

Interleukin-17F (IL-17F) exists in a disulfide-linked heterodimer that belongs to the IL-17 family. IL-17F is expressed in activated, but not resting, CD4+ T-cells and activated monocytes. Mouse and human IL-17F share 55% sequence identity.IL-17F has been shown to stimulate the production of several other cytokines, including IL-6, IL-8, and granulocyte colony-stimulating factor. IL-17F can regulate cartilage matrix turnover and stimulates PBMC and T-cell proliferation. IL-17F is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. Defects in IL-17F are the cause of familial candidiasis type 6 (CANDF6).

For Research Use Only