## Recombinant Mouse IL11RA/IL11R? Protein (His Tag)

Catalog Number: PKSM040940



Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Synonyms** AI314697;GP130;II-11ra;II11ra;II11ra2;NR1;RP23-167I12.2

Species Mouse

Expression Host HEK293 Cells
Sequence Met 1-Gln 367
Accession NP 001156873.1

Calculated Molecular Weight 39.2 kDa
Observed molecular weight 45-50 kDa
Tag C-His

**Bioactivity** Measured by its ability to bind mouse IL6ST in a functional ELISA.

#### **Properties**

**Purity** > 85 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 85 % as determined by reducing SDS-PAGE.

### **Background**

Interleukin 11 receptor, alpha subunit (IL11RA/IL-11RA) is a subunit of the interleukin 11 receptor which is a member of the hematopoietic cytokine receptor family. IL11RA/IL-11RA is expressed in a number of cell lines, including the myelogenous leukemia cell line K562, the megakaryocytic leukemia cell line Mo7E, the erythroleukemia cell line TF1, and the osteosarcoma cell lines, MG-63 and Saos-2. It is also expressed in normal and malignant prostate epithelial cell lines. Expression levels are increased in prostate carcinoma. This particular receptor is very similar to ciliary neurotrophic factor, since both contain an extracellular region with a 2-domain structure composed of an immunoglobulin-like domain and a cytokine receptor-like domain. Alternative splicing has been observed at this locus, and three variants encoding two

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Mouse IL11RA/IL11R? Protein (His Tag)

Catalog Number:PKSM040940



different isoforms have been identified. IL11RA/IL-11RA is a receptor for interleukin-11. The receptor systems for IL6, LIF, OSM, CNTF, IL11 and CT1 can utilize IL6ST for initiating signal transmission. Defects in IL11RA/IL-11RA are a cause of craniosynostosis and dental anomalies (CRSDA). CRSDA is a disorder characterized by craniosynostosis, maxillary hypoplasia, and dental anomalies, including malocclusion, delayed and ectopic tooth eruption, and/or supernumerary teeth. Some patients also display minor digit anomalies, such as syndactyly and/or clinodactyly.

#### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a>
 Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>