## **Elabscience**®

## Purified Anti-Human CD3 Antibody[UCHT1]

## catalog number: E-AB-F1230A

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description		
Reactivity	Human	
Immunogen	Recombinant Human CD3 protein	
Host	Mouse	
Isotype	Mouse IgG1, ĸ	
Clone	UCHT1	
Purification	>98%, Protein A/G purified	
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.	

 $2 \,\mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

Applications	<b>Recommended Dilution</b>

FCM

<u>Data</u>



Human peripheral blood lymphocytes were stained with 0.2 μg Purified Anti-Human CD3 Antibody[UCHT1](Right) and 0.2 μg Mouse IgG1, κ Isotype Control(Left), followed by PEconjugated Goat Anti-Mouse IgG Secondary Antibody

conjugated Goat Anti-Mouse igo secondary Antibody.	
Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles.
Shipping	Ice bag
Background	

CD3 $\epsilon$  is a 20 kD chain of the CD3/T cell receptor (TCR) complex, which is composed of two CD3 $\epsilon$ , one CD3 $\gamma$ , one CD3 $\delta$ , one CD3 $\zeta$  (CD247), and a T cell receptor ( $\alpha/\beta$  or  $\gamma/\delta$ ) heterodimer. It is found on all mature T lymphocytes, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.

## For Research Use Only