

A Reliable Research Partner in Life Science and Medicine

PE Anti-Mouse CD11c Antibody[N418]

Catalog Number: E-AB-F0991UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Host Armenian Hamster
Isotype Armenian Hamster IgG

Clone No. N418

Isotype Control PE Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells is 4.00 μ L values 1

in 100 µL volume].

<u>Data</u>



C57BL/6 murine splenocytes are stained with PE Anti-Mouse CD11c Antibody and APC Anti-Mouse MHC II (I-A/I-E) Antibody (Left). Splenocytes stained with APC Anti-Mouse MHC II (I-A/I-E) Antibody (Right) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion

receptor p150+95

Uniprot ID Q9QXH4

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 16411

CD11c is a 150 kD glycoprotein also known as α X integrin, CR4, and p150. CD11c forms a α X β 2 heterodimer with β 2 integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The α X β 2 integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen and CD54.