# Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410]

catalog number: E-AB-F1016A



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

**Host** Mouse


 $\begin{tabular}{lll} \textbf{Isotype} & Mouse IgGl, \kappa \\ \textbf{Clone} & MIAP410 \\ \textbf{Conjugation} & Unconjugated \\ \end{tabular}$ 

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

**Applications** Recommended Dilution

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

#### Data



Rat splenocytes were stained with 0.2  $\mu$ g Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410] (Right) and 0.2  $\mu$ g Mouse IgG1,  $\kappa$  Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

### **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** Ice bag

## Background

CD47, also known as Integrin-Associated Protein (IAP), is a membrane protein of about 50 kD with an IgV-like extracelluluar domain, a five membrane-spanning segment and a short terminal cytoplasmic region. It is widely expressed on many cell types and often associated with beta 3 integrins. It has been reported that CD47 functions as a self marker. Red cells lacking CD47 were rapidly cleared from the bloodstream by splenic macrophages. By binding to SIRP $\alpha$ , CD47 controls hemostatic innate immune functions, such as phagocytosis and cell trafficking.

### For Research Use Only