

# Recombinant Human Pleiotrophin/PTN/HB-GAM Protein



Catalog Number:PKSH031708

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	HARP;HBGF8;HBNF;NEGF1
<b>Species</b>	Human
<b>Expression Host</b>	Baculovirus-Insect Cells
<b>Sequence</b>	Met 1-Asp168
<b>Accession</b>	NP_002816.1
<b>Calculated Molecular Weight</b>	15.3 kDa
<b>Observed molecular weight</b>	19 kDa
<b>Tag</b>	None
<b>Bioactivity</b>	Immobilized human PTN at 10 µg/ml (100 µl/well) can bind rat SDC1-Fc, The EC50 of rat SDC1-Fc is 0.35-0.81 µg/ml.

## Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 1M NaCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 90 % as determined by reducing SDS-PAGE.

## Background

HB-GAM belongs to the pleiotrophin family. During embryonic and early postnatal development, HB-GAM is expressed in the central and peripheral nervous system and also in several non-neuronal tissues, notably lung, kidney, gut and bone. While in the adult central nervous system, it is expressed in an activity-dependent manner in the hippocampus where it can suppress long term potentiation induction. HB-GAM has a low expression in other areas of the adult brain, but it can be induced by ischemic insults, or targeted neuronal damaged in the entorhinal cortex or in the substantia nigra pars compacta. It is structurally related to midkine and retinoic acid induced heparin-binding protein and has a high affinity for

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

# Recombinant Human Pleiotrophin/PTN/HB-GAM Protein



Catalog Number:PKSH031708

heparin. HB-GAM binds anaplastic lymphoma kinase (ALK) which induces MAPK pathway activation, an important step in the anti-apoptotic signaling of PTN and regulation of cell proliferation. It also functions as a secreted growth factor and induces neurite outgrowth and which is mitogenic for fibroblasts, epithelial, and endothelial cells.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017