

## Recombinant Rat CD16a/FCGR3A Protein (His Tag)

**Catalog Number:** PKSR030403

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	HEK293 Cells-derived Rat CD16a/FCGR3A protein Met 1-Pro 200, with an C-terminal His
<b>Calculated MW</b>	21.7 kDa
<b>Observed MW</b>	28-33 kDa
<b>Accession</b>	XP_008767961.1
<b>Bio-activity</b>	Immobilized rat FCGR3A-His at 10 µg/ml (100 µl/well) can bind biotinylated human IgG1, The EC <sub>50</sub> of biotinylated human IgG1 is 59.0-139.0 ng/ml.

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

The Fc receptor with low affinity for IgG (FCGR3, or CD16) is encoded by 2 nearly identical genes, FCGR3A and FCGR3B, resulting in tissue-specific expression of alternative membrane-anchored isoforms. FCGR3A, it is also known as CD16a, encodes a transmembrane protein expressed on activated monocytes/macrophages, natural killer (NK) cells, and a subset of T cells.

CD16a / FCGR3A is a receptor expressed on NK cells that facilitates antibody dependent cellular cytotoxicity (ADCC) by binding to the Fc portion of various antibodies. CD16a / FCGR3A also has a broader function. CD16a / FCGR3A is directly involved in the lysis of some virus-infected cells and tumor cells by NK cells, independent of antibody binding. Cross-linking of CD16a / FCGR3A on NK cells resulted in increased intracellular Ca<sup>2+</sup> levels and a cascade of biochemical events similar to those activated by the T cell receptor. CD16a / FCGR3A on human NK cells is a lysis receptor that mediates the direct killing of some virus infected and tumor cells, independent of antibody ligation.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017