



A Reliable Research Partner in Life Science and Medicine

# **Monkey sCD14 Antibody Pair Set**

Catalog No. E-KAB-0663 Applications ELISA

Synonyms CD14 Antigen; Monocyte Differentiation Antigen CD14

## Kit components & Storage

Title	Specifications	Storage
Monkey sCD14 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Monkey sCD14 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0663)		
		Monkey sCD14 Capture Antibody	Monkey sCD14 Detection Antibody (Biotin)	
Immunogen	Immunogen	Recombinant Monkey sCD14 protein	Recombinant Monkey sCD14 protein	
Information	Swissprot	Q95M53		
Product details	Reactivity	Monkey	Monkey	
	Host	Mouse	Sheep	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Monkey sCD14 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Monkey sCD14 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Monkey sCD14 Capture	
Capture		Antibody	10
			8 II
			Optical Density
ELISA	1:1000-1:10000	Monkey sCD14 Detection	ojido o.1-
Detection		Antibody (Biotin)	2
			0.01 1 10 100 1000
			Monkey sCD14 Concentration (ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

CD14 is a 50-55 kDa glycosylphosphatidylinositol-anchored glycoprotein. CD14 is preferentially expressed on monocytes and macrophages. CD14 acts as a co-receptor (along with TLR4 and MD-2) for bacterial liposaccharides (LPS). It plays a major role in the inflammatory response of monocytes to LPS.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>