# Recombinant Human Protocadherin-10/PCDH10 Protein (Fc Tag)

**Catalog Number: PKSH032979**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Species** | Human |
| **Source** | HEK293 Cells-derived Human Protocadherin-10;PCDH10 protein Gln19-Thr715, with an C-terminal Fc |
| **Calculated MW** | 102.4 kDa |
| **Observed MW** | 125-165 kDa |
| **Accession** | Q9P2E7 |
| **Bio-activity** | Not validated for activity |

## Properties

| | |
|---|---|
| **Purity** | > 75 % as determined by reducing SDS-PAGE. |
| **Endotoxin** | < 1.0 EU per μg of the protein as determined by the LAL method. |
| **Storage** | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 ℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months. |
| **Shipping** | This product is provided as lyophilized powder which is shipped with ice packs. |
| **Formulation** | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.<br>Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| **Reconstitution** | Please refer to the printed manual for detailed information. |

## Data




> 75 % as determined by reducing SDS-PAGE.

## Background

Protocadherin-10 (PCDH10) is a single-pass type I membrane protein that contains six extracellular cadherin domains; one transmembrane domain and one cytoplasmic tail differing from those of the classical cadherins. As a potential calcium-dependent cell-adhesion neuronal receptor; it may plays a role in the establishment and function of specific cell-cell connections in the brain. PCDH10 moderately expressed in all regions of the brain examined; as well as in testis and ovary; and low expression in all other tested tissues.