## Recombinant Rat ST14/Matriptase protein (His Tag)

## Catalog Number: PDER100225

Note: Centrifuge before opening to ensure complete recovery of vial contents.

reconstituted samples are stable at < -20°C for 3 months.	Description			
Calculated MW29.0 kDaObserved MW34 kDaAccessionQ9JJI7Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Species	Rat		
Observed MW34 kDaAccessionQ9JJ17Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Source	E.coli-derived Rat ST14 protein Lys 591-Val855, with an N-terminal His		
AccessionQ9JJI7Bio-activityNot validated for activityPropertiesPurity>95% as determined by reducing SDS-PAGE.Findotoxin< 10 EU/mg of the protein as determined by the LAL method	Calculated MW	29.0 kDa		
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Observed MW	34 kDa		
Properties   Purity > 95% as determined by reducing SDS-PAGE.   Endotoxin < 10 EU/mg of the protein as determined by the LAL method   Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20   °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots reconstituted samples are stable at < -20°C for 3 months.   Shipping This product is provided as lyophilized powder which is shipped with ice packs.   Formulation Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%	Accession	Q9JJI7		
Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	<b>Bio-activity</b>	Not validated for activity		
Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Properties			
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots reconstituted samples are stable at < -20°C for 3 months.	Purity	>95% as determined by reducing SDS-PAGE.		
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 10 EU/mg of the protein as determined by the LAL method		
reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -2		
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Mannitol	Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%		
triumnor.		Mannitol.		
<b>Reconstitution</b> It is recommended that sterile water be added to the vial to prepare a stock solut	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
0.5 mg/mL. Concentration is measured by UV-Vis.		0.5 mg/mL. Concentration is measured by UV-Vis.		

## Data

KDa	М	R
180 135	-	
100		-
75	-	-
65	-	-
45	-	-
35		-
25		

> 95 % as determined by reducing SDS-PAGE.

## Background

Degrades extracellular matrix. Proposed to play a role in breast cancer invasion and metastasis. Exhibits trypsin-like activity as defined by cleavage of synthetic substrates with Arg or Lys as the P1 site.Defects in ST14 are a cause of ichthyosis autosomal recessive with hypotrichosis (ARIH). ARIH is a skin disorder characterized by congenital ichthyosis associated with the presence of less than the normal amount of hair.