

Recombinant Human ERN1/IRE1 Protein (aa 465-977)

Catalog Number: PKSH030975

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human ERN1/IRE1 protein Pro 465-Leu 977
Calculated MW	58.3 kDa
Observed MW	65 kDa
Accession	O75460-1
Bio-activity	1. Kinase activity untested 2. Measured by its nuclease activity to cleave Xbp1 single stem-loop mini-substrate.

Properties

Purity	> 80 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 80 % as determined by reducing SDS-PAGE.

Background

Trypsin-3; also known as Trypsin III; brain trypsinogen; Serine protease 3 and PRSS3; is a secreted protein which belongs to the peptidase S1 family. Trypsin-3 / PRSS3 is expressed in pancreas and brain. It contains one peptidase S1 domain. Trypsin-3 / PRSS3 can degrade intrapancreatic trypsin inhibitors that protect against CP. Genetic variants that cause higher mesotrypsin activity might increase the risk for chronic pancreatitis (CP). A sustained imbalance of pancreatic proteases and their inhibitors seems to be important for the development of CP. The trypsin inhibitor-degrading activity qualified PRSS3 as a candidate for a novel CP susceptibility gene. Trypsin-3 / PRSS3 has been implicated as a putative tumor suppressor gene due to its loss of expression; which is correlated with promoter hypermethylation; in esophageal squamous cell carcinoma and gastric adenocarcinoma.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017