# **RAF1** Polyclonal Antibody

catalog number: E-AB-60020



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human RAF1 (NP_002871.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

oplications	Recommended Dilution
В	1:500-1:2000
С	1:50-1:200
	1:50-1:200
B C	1:50-1:2000 1:50-1:200 1:50-1:200

#### Data



Western blot analysis of extracts of various cell lines using RAF1 Polyclonal Antibody at dilution of 1:1000.

## Observed-MV:73 kDa Calculated-MV:73 kDa/75 kDa



nistry of paraffin-embedde

Immunohistochemistry of paraffin-embedded Rat liver using RAF1 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Rat heart using RAF1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunofluorescence analysis of U2OS cells using RAF1 Polyclonal Antibody

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

### Background

## For Research Use Only

# **RAF1** Polyclonal Antibody

### catalog number: E-AB-60020



This gene is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in this gene are associated with Noonan syndrome 5 and LEOPARD syndrome 2.

### For Research Use Only