

A Reliable Research Partner in Life Science and Medicine

## APC Anti-Mouse CD105 Antibody[MJ7/18]

Catalog Number: E-AB-F1233E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse Host Rat

**Isotype** Rat IgG2a, κ **Clone No.** MJ7/18

Isotype Control APC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832E]

Conjugation APC

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

#### Applications Recommended usage

FCM Each lot of

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Data**



Mouse endothelial cells (bEnd.3) are stained with APC Anti-Mouse CD105 Antibody (filled gray histogram).APC Mouse IgG2a, κ Isotype Control stained cells are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

Rev. V1.5

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names END;Endoglin;Eng

 Uniprot ID
 Q63961

 Gene ID
 13805

### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

#### **Background**

CD105 is a 90 kD homodimeric type I integral membrane glycoprotein, also known as endoglin. It is expressed on endothelial cells (especially on angiogenic endothelial cells) and upregulated by hypoxia, activated monocytes, macrophages, bone marrow stromal cells, and some cytotrophoblasts. CD105 is a receptor for TGF- $\beta$ 1, TGF- $\beta$ 3 and modulates TGF- $\beta$  signaling by interacting with TGF- $\beta$  receptors I and/or II. CD105 also binds other growth factors such as actvin A, BMP-2, and BMP-7. CD105 has been show to be a useful marker for identifying proliferating endothelium involved in tumor angiogenesis and can be used for tumor imaging and prognosis, and has therapeutic potential for some solid tumors and other angiogenic diseases.

Fax: 1-832-243-6017