

A Reliable Research Partner in Life Science and Medicine

# **RIDA Polyclonal Antibody**

catalog number: E-AB-53037

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human; Mouse; Rat

Immunogen Fusion protein of human RIDA

Host Rabbit Isotype IgG


**Purification** Antigen affinity purification

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


### **Applications** Recommended Dilution

**WB** 1:500-1:2000 **IHC** 1:50-1:200

#### Data




Western blot analysis of Mouse liver tissue and Mouse kidney tissue lysates using RIDA Polyclonal Antibody at dilution of 1:650



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using RIDA Polyclonal Antibody at dilution of  $1:70(\times 200)$ 

## Observed-MW:Refer to figures Calculated-MW:14 kDa



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using RIDA Polyclonal Antibody at dilution of 1:70(×200)

### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

# Background

### For Research Use Only

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

Catalyzes the hydrolytic deamination of enamine/imine intermediates that form during the course of normal metabolism. May facilitate the release of ammonia from these potentially toxic reactive metabolites, reducing their impact on cellular components. It may act on enamine/imine intermediates formed by several types of pyridoxal-5'-phosphate-dependent dehydratases including L-threonine dehydratase. May also function as an endoribonuclease, cleaving mRNA phosphodiester bonds of single-stranded RNA (By similarity). Thereby, may inhibit protein translation (PubMed: 8973653).

Fax: 1-832-243-6017