

PerCP/Cyanine 5.5 Anti-Mouse/Rat CD29 Antibody[HMβ1-1]

Catalog Number: E-AB-F1309J

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|--------------------------------|---|
| Reactivity | Mouse,Rat |
| Host | Armenian Hamster |
| Isotype | Armenian Hamster IgG |
| Clone No. | HMβ1-1 |
| Isotype Control | PerCP/Cyanine5.5 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852J] |
| Conjugation | PerCP/Cyanine 5.5 |
| Conjugation Information | PerCP/Cyanine 5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter). |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. |

Applications

Recommended usage

| | |
|------------|---|
| FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |
|------------|---|

Data



C57BL/6 murine splenocytes are stained with PerCP/Cyanine 5.5 Anti-Mouse/Rat CD29 Antibody (filled gray histogram) or PerCP/Cyanine 5.5 Armenian Hamster IgG Isotype Control (empty black histogram).

Preparation & Storage

| | |
|-----------------|---|
| Storage | Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag |

Antigen Information

| | |
|------------------------|---|
| Alternate Names | integrin β1;VLA-β chain;β1 integrin;GP11a;ITGB1 |
| Uniprot ID | P09055; P49134 |
| Gene ID | 16412; 24511 |

For Research Use Only

Background

CD29 is a 130 kD protein, also known as integrin $\beta 1$, VLA- β chain, or GPIIa. It is a member of the integrin family, expressed broadly on leukocytes, endothelial cells, smooth muscle, and epithelial cells. In association with CD49a-f, CD29 forms the VLA-1 through VLA-6 complexes, respectively. It plays an important role in cell-cell or cell-matrix interaction. The HM β 1-1 antibody reacts with both mouse and rat CD29. It is able to block cell adhesion and inhibit T cell proliferation.