

Recombinant MAP2/METAP2 Monoclonal Antibody

catalog number: **AN300529P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Mouse
Immunogen	Recombinant Mouse MAP2/METAP2 protein
Host	Rabbit
Isotype	IgG
Clone	7A3
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications Recommended Dilution


WB	1:500-1:2000
IP	0.5-2 µL/mg of lysate

Data



Western Blot with METAP2 Monoclonal Antibody at dilution of 1:500. Lane A: HepG2 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:65 kDa
Calculated-MW:52 kDa



Immunoprecipitation analysis using 2 µL anti-Mouse METAP2 Monoclonal Antibody and 15 µL of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using METAP2 Monoclonal Antibody at a dilution of 1:200. Lane A:0.5 mg HepG2 Whole Cell Lysate

Observed-MW:65 kDa
Calculated-MW:52 kDa

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

METAP2 (Methionine aminopeptidase 2), also known as MAP2 is a protein that belongs to the peptidase M24A family. MAP2 binds 2 cobalt or manganese ions and contains approximately 12 O-linked N-acetylglucosamine (GlcNAc) residues. It is found in all organisms and is especially important because of its critical role in tissue repair and protein degradation. The catalytic activity of human MAP2 toward Met-Val peptides is consistently two orders of magnitude higher than that of METAP1, suggesting that it is responsible for processing proteins containing N-terminal Met-Val and Met-Thr sequences in vivo. This protein functions both by protecting the alpha subunit of eukaryotic initiation factor 2 from inhibitory phosphorylation and by removing the amino-terminal methionine residue from nascent protein. MAP2 protects eukaryotic initiation factor EIF2S1 from translation-inhibiting phosphorylation by inhibitory kinases such as EIF2AK2/PKR and EIF2AK1/HCR. It also plays a critical role in the regulation of protein synthesis.