

Recombinant WSTF Monoclonal Antibody

catalog number: AN301948L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Rat;
Immunogen	Recombinant human WSTF fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A664
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

WB	1:1000
-----------	--------

Data



Western Blot with WSTF Monoclonal Antibody at dilution of 1:1000. Lane 1: T47D, Lane 2: MCF-7, Lane 3: PC-12, Lane 4: SH-SY5Y

Observed-MW:171 kDa

Calculated-MW:171 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The human WSTF gene is located within the common Williams Syndrome (WS) deletion area at chromosome 7q11.23. Several WSTF gene products have been detected with little difference in length of polypeptides. Functional motifs identified by sequence-homology searches include a PHD-type zinc finger motif followed by a bromodomain. Both motifs are found in many transcription factors, suggesting that WSTF may function as a transcription factor. A *Drosophila* gene (*acf1*) was cloned, which encodes two forms of Acf1 proteins with molecular weight 170 kDa and 185 kDa, respectively. It was demonstrated that Acf1 is structurally related to the human WSTF gene. Acf1 forms a complex with another protein, ISWI, and functions in the ATP-dependent catalysis of chromatin assembly.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0