

A Reliable Research Partner in Life Science and Medicine

PerCP/Cyanine5.5 Anti-Human CD69 Antibody[FN50]

Catalog Number: E-AB-F1138J

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ

Clone No. FN50

Isotype Control PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792J]

Conjugation PerCP/Cyanine 5.5

Conjugation Information PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected

using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



PMA and ionomycin-stimulated (4h) Jurkat cells are stained with PerCP/Cyanine5.5 Anti-Human CD69 Antibody (filled gray histogram) or Mouse IgG1 Isotype Control PerCP/Cyanine5.5 (empty black histogram).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8 $^{\circ}\text{C}$ for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names AlM; Activation inducer molecule; EA1; Early activation antigen CD69; MLR-3

 Uniprot ID
 Q07108

 Gene ID
 969

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience[®]

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD69 is a 27-33 kD type II transmembrane protein also known as activation inducer molecule (AIM), very early activation antigen (VEA), and MLR3. It is a member of the C-type lectin family, expressed as a disulfide-linked homodimer. Other members of this receptor family include NKG2, NKR-P1 CD94, and Ly49. CD69 is transiently expressed on activated leukocytes including T cells, thymocytes, B cells, NK cells, neutrophils, and eosinophils. CD69 is constitutively expressed by a subset of medullary mature thymocytes, platelets, mantle B cells, and certain CD4+ T cells in germinal centers of normal lymph nodes. CD69 is involved in early events of lymphocyte, monocyte, and platelet activation, and has a functional role in redirected lysis mediated by activated NK cells.

Fax: 1-832-243-6017