

Recombinant Human Contactin 5/CNTN5 Protein (His Tag)



Catalog Number:PKSH031559

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	HNB-2s;NB-2
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Gln 1059
Accession	NP_055176.1
Calculated Molecular Weight	116 kDa
Observed molecular weight	130-140 kDa
Tag	C-His
Bioactivity	Measured by the ability of the immobilized protein to support the adhesion of C6 Rat brain glial cells. When 5×10^4 cells/well are added to CNTN5 coated plates (0.8 $\mu\text{g}/\text{ml}$ and 100 $\mu\text{l}/\text{well}$), approximately 30%-50% will adhere specifically after 60 minutes at 37°C.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed mainly in the nervous system. The subgroup consists of six members: Contactin-1, Contactin-2(TAG-1), Contactin-3(BIG-1), BIG-2, Contactin-5(NB-2) and NB-3. Since their identification in the late 1980s, Contactin-1 and Contactin-2 have been studied extensively. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human Contactin 5/CNTN5 Protein (His Tag)



Catalog Number:PKSH031559

study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. In contrast, the function of the other four members of this subgroup remained unknown until recently. Contactin-5, also known as NB-2, is one of the neural recognition molecules in the contactin subgroup. Contactin-5 is expressed in brain and kidney and at very low level in placenta. In brain, Contactin-5 is highly expressed in the occipital lobe, amygdala, cerebral cortex, frontal lobe, thalamus and temporal lobe. Mice deficient in the Contactin-5 gene exhibit aberrant responses to acoustic stimuli. Contactin-5 may play a role in maturation of glutamatergic synapses in the brainstem during the final stages of auditory development. Contactin-5 gene may contribute to human neurological disorders.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017