INHA Polyclonal Antibody

catalog number: E-AB-14159





Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Recombinant protein of human INHA |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Pacammandad Dilution |

| Applications | Recommended Dilution |
|--------------|-----------------------------|
| IHC | 1:150-1:500 |

Data





Immunohistochemistry of paraffin-embedded Human tonsil tissue using INHA Polyclonal Antibody at dilution 1:150 Immunohistochemistry of paraffin-embedded Human renal cancer tissue using INHA Polyclonal Antibody at dilution

1:150

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |

Background

This gene encodes the alpha subunit of inhibins A and B protein complexes. These complexes negatively regulate follicle stimulating hormone secretion from the pituitary gland. Inhibins have also been implicated in regulating numerous cellular processes including cell proliferation, apoptosis, immune response and hormone secretion.