## Recombinant Mouse IL-25 protein(His Tag)

Catalog Number: PKSM041471



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	EDF;BCDFII;TRF
Species	Mouse
Expression Host	E.coli
Sequence	Val 17-Ala 169
Accession	NP_542767.1
Calculated Molecular Weight	18.4 kDa
Observed molecular weight	17 kDa
Tag	C-His
Bioactivity	Measured by its ability to induce CXCL1 secretion in HT-29 cells. The $ED_{50}$ for this effect is < 1 ng/mL
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per $\mu$ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20 mM sodium citrate, 0.2 M NaCl, pH 4.5. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	
kDa 75- 63- 48- 35- 25- 17- 11-	

> 98 % as determined by reducing SDS-PAGE.

## Background

Interleukin-25 (IL-25) is a cytokine that shares sequence similarity with interleukin 17. This cytokine can induce NFkappaB activation, and stimulate the production of interleukin 8. Both this cytokine and interleukin 17B are ligands for the cytokine receptor IL17BR. IL-25 is a member of the IL-17 family of cytokines. However, unlike the other members of this family, IL-25 promotes T helper (Th) 2 responses. IL-25 also regulates the development of autoimmune inflammation mediated by IL-17–producing T cells. IL-25 and IL-17, being members of the same cytokine family, play opposing roles in the pathogenesis of organ-specific autoimmunity. IL-25 promotes cell expansion and Th2 cytokine

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number:PKSM041471



production when Th2 central memory cells are stimulated with thymic stromal lymphopoietin (TSLP)–activated dendritic cells (DCs), homeostatic cytokines, or T cell receptor for antigen triggering. Elevated expression of IL-25 and IL-25R transcripts was observed in asthmatic lung tissues and atopic dermatitis skin lesions, linking their possible roles with exacerbated allergic disorders. A plausible explanation that IL-25 produced by innate effector eosinophils and basophils may augment the allergic inflammation by enhancing the maintenance and functions of adaptive Th2 memory cells had been provided.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com