Recombinant Human IL2Ra/CD25 Protein (Fc Tag)

Catalog Number: PKSH032570



Note: Centrifuge before opening to ensure complete recovery of vial contents.

	escrip				7.0		
	00		TO T	m	TT.		т
v		ж			ш	w	٠

 Species
 Human

 Mol_Mass
 48.9 kDa

 Accession
 P01589

Bio-activity Loaded Human IL-2-His on HIS1K Biosensor, can bind Human IL-2RA-Fc with an

affinity constant of 3.06 nM as determined in BLI assay.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-2 receptor subunit alpha (II.2RA) is a single-pass type I membrane protein; contains 2 Sushi (CCP/SCR) domains. The interleukin 2 (II.2) receptor alpha (II.2RA) and beta (II.2RB) chains; together with the common gamma chain (II.2RG); constitute the high-affinity II.2 receptor. Homodimeric alpha chains (II.2RA) result in low-affinity receptor; while homodimeric beta (II.2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein; soluble II.2RA has been isolated and determined to result from extracellular proteolyisis.

For Research Use Only