

A Reliable Research Partner in Life Science and Medicine

PerCP Anti-Mouse CD5 Antibody[53-7.3]

Catalog Number: E-AB-F1185UF

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ **Clone No.** 53-7.3

Isotype Control PerCP Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833F]

Conjugation PerCP

Conjugation Information PerCP is designed to be excited by the blue laser (488 nm) and detected using an optical

filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].

Data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD45R/B220 Antibody and PerCP Anti-Mouse CD5 Antibody (Left). Splenocytes are stained with FITC Anti-Mouse CD45R/B220 Antibody and PerCP Rat IgG2a, κ Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD5;Cd5;Ly-1;Lymphocyte antigen 1;Lyt-1

 Uniprot ID
 P13379

 Gene ID
 12507

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

CD5 is a 67 kD protein, also known as Lyt-1, Ly-1, T1, Tp67, or Ly-12. It is a member of the scavenger receptor cysteine-rich protein superfamily (SRCR) and primarily expressed on thymocytes, T cells, and B-1 cells. Although mature α/β T cells express high levels of CD5, very few γ/δ T cells express this antigen. The interaction of CD5 with CD72, gp35-37, TCR, or BCR is involved in T and B cell activation.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com