Recombinant Mouse Adiponectin/AdipoQ Protein (E.coli, His Tag)

Catalog Number: PKSM040960



Note: Centrifuge before opening to ensure complete recovery of vial contents. Description Species Mouse Mol Mass 27.2 kDa Accession O60994 Not validated for activity **Bio-activity Properties** > 90 % as determined by reducing SDS-PAGE. Purity < 1.0 EU per µg of the protein as determined by the LAL method. Endotoxin Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Shipping Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Formulation Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. Reconstitution Please refer to the printed manual for detailed information. Data

	kDa	MKence	R
	120 90 60	1200	chence
	40	Ele	10-2
	30		cience
	20		
	14	-	

> 90 % as determined by reducing SDS-PAGE.

Background

Adiponectin is a secreted protein. It is synthesized exclusively by adipocytes and secreted into plasma. Adiponectin is an important adipokine that is involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Adiponectin Stimulates AMPK phosphorylation and activates in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Adiponectin also antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. It inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. Adiponectin may play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex: LMW, MMW or HMW.

For Research Use Only