

PE/Cyanine7 Anti-Human Myeloperoxidase Antibody[1B10]

Catalog Number: AN00925H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	1B10
Isotype Control	PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792H]
Conjugation	PE/Cyanine 7
Conjugation Information	PE/Cyanine 7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

Data



Intracellular staining of normal human peripheral blood cells with PE/Cyanine7 Anti-Human Myeloperoxidase Antibody[1B10] (filled gray histogram) or PE/Cyanine7 Mouse IgG1, κ Isotype Control (empty black histogram). Cells in the granulocytes gate were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protect from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	MPO
Uniprot ID	P05164

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017
Rev. V1.3

Gene ID	4353
Background	<p>Myeloperoxidase (MPO) is a heterotetrameric protein consisting of two 60 kD heavy units and two 12 kD light units. A lysosomal enzyme, MPO is able to catalyze the production of hypochlorous acid, a potent microbicidal agent, from hydrogen peroxide and chloride anion during the neutrophil respiratory burst. MPO is a major enzyme involved in the inflammatory responses of polymorphonuclear leucocytes. MPO is localized to the azurophilic granules of mature granulocytes and monocytes and is also expressed in some acute myeloid leukemia cells.</p>

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3