Elabscience®


Elab Fluor[®] Violet 450 Anti-Human CD3 Antibody[UCHT1]

Catalog Number: E-AB-F1230Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|-------------------------|--|
| Reactivity | Human |
| Host | Mouse |
| lsotype | Mouse IgG1, κ |
| Clone No. | UCHT1 |
| Isotype Control | Elab Fluor [®] Violet 450 Mouse IgG1, к Isotype Control[MOPC-21] [Product E-AB- F09792Q] |
| Conjugation | Elab Fluor [®] Violet 450 |
| Conjugation Information | Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a $450/45$ nm bandpass filter). |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA. |
| Applications | Recommended usage |
| FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |

Data



Human peripheral blood lymphocytes are stained with Elab

Fluor[®] Violet 450 Anti-Human CD3 Antibody (filled gray histogram).Unstained splenocytes (empty black histogram) are used as control.

| Preparation & Storag | ge |
|----------------------|---|
| Storage | Keep as concentrated solution. |
| | This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag |
| Antigen Information | |
| Alternate Names | CD3E;CD3e;T-cell surface antigen T3/Leu-4 epsilon chain;T-cell surface glycoprotein CD3 epsilon chain;T3E |
| Uniprot ID | P07766 |
| Gene ID | 915 |
| For Research Use | Only |
| Tal: 400 000 2100 | |

Elabscience®

Background

Elabscience Biotechnology Co., Ltd. A Reliable Research Partner in Life Science and Medicine

CD3 ϵ is a 20 kD chain of the CD3/T cell receptor (TCR) complex, which is composed of two CD3 ϵ , one CD3 γ , one CD3 δ , one CD3 ζ (CD247), and a T cell receptor (α/β or γ/δ) heterodimer. It is found on all mature T lymphocytes, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.