A Reliable Research Partner in Life Science and Medicine

PE/Cyanine5 Anti-Mouse CD183/CXCR3 Antibody[CXCR3-173]

Catalog Number: E-AB-F1114G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Host Armenian Hamster Isotype Armenian Hamster IgG

Clone No. CXCR3-173

[Product E-AB-F09852G] Isotype Control

PE/Cyanine 5 Conjugation

Conjugation Information PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm

(e.g., a 690/50 nm bandpass filter).

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



C57BL/6 murine splenocytes are stained with PE/Cyanine5

Anti-Mouse CD183/CXCR3 Antibody and Elab Fluor® 488 Anti-Mouse CD3 Antibody (Left). Splenocytes stained with

Elab Fluor® 488 Anti-Mouse CD3 Antibody Antibody (Right) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names C-X-C chemokine receptor type 3;CD183/CXCR3;CXC-R3;CXCR-3;Cxcr3;IP-10

receptor;Interferon-inducible protein 10 receptor

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Uniprot ID Gene ID Background O88410 12766

CD183/CXCR3, also known as CXCR3, is a member of the C-X-C chemokine family, characterized by a pair of cysteine residues separated by a single amino acid. CXCR3 is a 38 kD seven pass transmembrane receptor coupled to G-protein. It mediates Ca2 + mobilization and chemotaxis in response to C-X-C chemokines, such as IP10 (CXCL10), MIG (CXCL9), I-TAC (CXCL11) and PF4 (CXCL4). CXCR3 is expressed primarily on activiated T lymphocytes, NK cells, and some epithelial cells and endothelial cells. It is not expressed on B cells, monocytes or granulocytes.

Fax: 1-832-243-6017