



Recombinant CD33 Monoclonal Antibody

catalog number: AN301475L

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description	
Reactivity	Human
Immunogen	Recombinant human CD33 fragment
Host	Rabbit
Isotype	IgG, κ
Clone	C351098
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
IHC	1:200-1:1000

Data	
	
Immunohistochemistry of paraffin-embedded Human lymphoma using CD33 Monoclonal Antibody at dilution of 1:1000.	Immunohistochemistry of paraffin-embedded Human spleen using CD33 Monoclonal Antibody at dilution of 1:1000.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background	
Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state. Upon engagement of ligands such as C1q or sialylated glycoproteins, two immunoreceptor tyrosine-based inhibitory motifs (ITIMs) located in CD33 cytoplasmic tail are phosphorylated by Src-like kinases such as LCK. These phosphorylations provide docking sites for the recruitment and activation of protein-tyrosine phosphatases PTPN6/SHP-1 and PTPN11/SHP-2. In turn, these phosphatases regulate downstream pathways through dephosphorylation of signaling molecules.	

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1