

## Purified Anti-Human CD11b Antibody[HI11b]

**catalog number: E-AB-F1317A**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human CD11b protein
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG2b, $\kappa$
<b>Clone</b>	HI11b
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Applications

### Recommended Dilution

<b>FCM</b>	2 $\mu\text{g/mL}$ ( $0.5 \times 10^6$ - $1 \times 10^6$ cells)
------------	---

### Data



Human peripheral blood granulocytes cell were stained with 0.2  $\mu\text{g}$  Purified Anti-Human CD11b Antibody[HI11b] (Right) and 0.2  $\mu\text{g}$  Mouse IgG2b,  $\kappa$  Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

Human ITGAM (Integrin alpha M subunit) is a 127kDa (predicted) glycoprotein, a member of the Integrin family of proteins. The Integrin family proteins are heterodimeric transmembrane receptors composed of an alpha and a beta subunit. The Integrin alpha M subunit, also known as MAC-1 alpha subunit or CD11b, combines with the Integrin beta 2 subunit (CD18) to form the non-covalent heterodimer Integrin alpha M/ beta 2, also known as MAC-1 and complement receptor type 3 (CR3).

### For Research Use Only