



A Reliable Research Partner in Life Science and Medicine

MAPKAPK3/MK3 Monoclonal Antibody

catalog number: AN200072P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human MAPKAPK3/MK3 Protein

HostMouseIsotypeIgG2aClone4B6PurificationProtein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

IHC-P 1:50-1:200

Data



Immunohistochemistry of paraffin-embedded human skeletal muscle using MAPKAPK3/MK3 Monoclonal Antibody at dilution of 1:60.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

This gene encodes a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)- activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and repress the activity of E44, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and cell differentiation. Alternate splicing results in multiple transcript variants that encode the same protein.