IL-6/IFN-β2/BSF-2, Human, Recombinant

Cat. No. : PCK085

General Information

Synonyms	Interleukin-6;IL-6;B-Cell Stimulatory Factor 2;BSF-2;CTL Differentiation Factor;CDF; Hybridoma Growth Factor;Interferon Beta-2;IFN-Beta-2;IL6;IFNB2
Species	Human
Expression host	E.coli
Sequence	Val30-Met212
Accession	P05231
Mol mass	20.9 KDa
Expiration date	12 months
Bio activity	Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 20-100 pg/mL.


Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 5% Sucorse, 4% Mannitol, 0.05% PS-80, pH7.4.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


Background

Cytokines of the IL6/GCSF/MGF family are glycoproteins of about 170 to 180 amino acid residues that contain four conserved cysteine residues involved in two disulfide bonds. They have a compact, globular fold (similar to other interleukins), stabilized by the 2 disulfide bonds. One half of the structure is dominated by a 4 alpha-helix bundle with a left-handed twist; the helices are anti-parallel, with 2 overhand connections, which fall into a 2-stranded anti-parallel beta-sheet. The fourth alpha helix is important to the biological activity of the molecule. Interleukin-6 (IL-6) is an important proinflammatory and immunoregulatory cytokine expressed by various cells. Interleukin-6 has been shown to inhibit the growth of early stage and to promote the proliferation of advanced stage melanoma cells in vitro.

SDS-PAGE



Bio activity



Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 20-100 pg/mL.

Procell system®