

Purified Anti-Human CD227 Antibody[AR20.5]


catalog number: AN007620P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human CD227 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	AR20.5
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	PBS, pH 7.2. Contains 0.05% Proclin300.

Applications	Recommended Dilution
FCM	2 μ g/mL (1×10^5 - 5×10^5 cells)

Data



MCF-7 were stained with 0.2 μ g Purified Anti-Human CD227 Antibody[AR20.5] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

MUC-1 (Mucin-1; also PEM, PEMT, Episialin, tumor-associated mucin, and carcinoma associated mucin) is a 240-450 kDa type I transmembrane glycoprotein. Human MUC-1 is 1255 amino acids (aa) in length and contains a 23 aa signal sequence, a 1135 aa extracellular domain (ECD), a 23 aa transmembrane segment, and a 74 aa cytoplasmic tail. The ECD consists of degenerate tandem repeats and a tandem repeat region, which makes up the major portion of the protein. Potential O-glycosylation sites (serines and threonines) make up more than one-fourth of the amino acids. Splicing variants produce ten isoforms for human MUC-1. Human MUC-1 is 28% aa identical to Mouse MUC-1.