

Recombinant IL-1RA/IL1RN Monoclonal Antibody

catalog number: AN300077P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description



Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide corresponding to the center region of the human IL1Ra / IL1RN.
Host	Rabbit
Isotype	IgG
Clone	C110059
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications

Recommended Dilution

IHC	1:200-1:1000
WB	1:2000-1:10000
IF	1:200-1:1000
ELISA	1:5000-1:20000
IP	1:50-1:200

Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-IL1RA antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: Hela
Lane 2: K562 Lane 3: RAW264.7 Lane 4: Rat spleen
Predicted band size: 17kDa Observed band size: 20kDa

Human esophagus was stained with anti-IL1RA Rabbit antibody

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.6

disease:Genetic variation in IL1RN is associated with susceptibility to diabetic nephropathy [MIM:612628]; also called susceptibility to microvascular complications of diabetes type 4 (MVCD4) . Diabetic nephropathy is a kidney disease and resultant kidney function impairment due to the long standing effects of diabetes on the microvasculature (glomerulus) of the kidney. Features include increased urine protein and declining kidney function. , function:Inhibits the activity of IL-1 by binding to its receptor. Has no IL-1 like activity. ,online information:Interleukin-1 entry ,online information:The Singapore human mutation and polymorphism database ,pharmaceutical:Available under the name Kineret (Amgen) . Used for the reduction in signs and symptoms and slowing the progression of structural damage in moderately to severely active rheumatoid arthritis. ,similarity:Belongs to the IL-1 family. ,tissue specificity:The intracellular form of IL1RN is predominantly expressed in epithelial cells. ,