

Pepsinogen C/PGC Monoclonal Antibody

catalog number: AN200124P

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human
Immunogen	Recombinant Human Pepsinogen C / PGC protein
Host	Mouse
Isotype	Mouse IgG1
Clone	A972
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications

Recommended Dilution

IHC-P	1:50-1:200
-------	------------

Data



Immunohistochemistry of paraffin-embedded human stomach using Pepsinogen C / PGC Monoclonal Antibody at dilution of 1:60.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

Pepsinogen II is a zymogen secreted by the chief cells of the gastric glands and converted into pepsin in the presence of gastric acid or of pepsin itself. Among endoscopy clinic patients, PGII is a sensitive marker for extension of nonatrophic gastritis toward the corpus. PGI is a stable biomarker in assessment of fundic atrophy and has similar accuracy to PGI/PGII ratio among populations with prevalent nonatrophic pangastritis. Research also showed that decreased PG I level and PG I/II ratio are risk factors for gastric cancer. Combined use of serum PG I level and PG I/II ratio may help the early diagnosis of gastric cancer.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2