Elabscience®

LATS1 Polyclonal Antibody

catalog number: E-AB-93222

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | |
|--|---------------------------|---|
| Reactivity | Human;Mouse;Rat | |
| Immunogen | A synthetic peptide of hu | uman LATS1 |
| Host | Rabbit | |
| Isotype | IgG | |
| Purification | Affinity purification | |
| Buffer | Phosphate buffered solut | tion, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilu | tion |
| WB | 1:500-1:2000 | |
| IHC | 1:50-1:200 | |
| Data | | |
| | 250kDaLATS1 100kDa | |
| Western blot englysis | 70kDa- | Immunchistechemistry of pareffin embedded reteningle and |
| LATEL D. L. al. and the destination of the second strain second strain strain second s | | |
| LATSI Polycional Antibody at 1:1000 dilution. using | | using LAISI Polyclonal Antibody at dilution of 1:100 (40x |

Observed-MW:150 kDa Calculated-MW:76 kDa/126 kDa using LATS1 Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack,upon receipt,store it immediately at the |
| | temperature recommended. |

Background

The protein encoded by this gene is a putative serine/threonine kinase that localizes to the mitotic apparatus and complexes with cell cycle controller CDC2 kinase in early mitosis. The protein is phosphorylated in a cell-cycle dependent manner, with late prophase phosphorylation remaining through metaphase. The N-terminal region of the protein binds CDC2 to form a complex showing reduced H1 histone kinase activity, indicating a role as a negative regulator of CDC2/cyclin A. In addition, the C-terminal kinase domain binds to its own N-terminal region, suggesting potential negative regulation through interference with complex formation via intramolecular binding. Biochemical and genetic data suggest a role as a tumor suppressor. This is supported by studies in knockout mice showing development of soft-tissue sarcomas, ovarian stromal cell tumors and a high sensitivity to carcinogenic treatments.

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017