Elabscience®

Human sCD163 Antibody Pair Set

Catalog No.	E-KAB-0207	Applications	ELISA
Synonyms	M130, MM130		

Kit components & Storage

Title	Specifications	Storage
Human sCD163 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Human sCD163 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0207)	
		Human sCD163 Capture Antibody	Human sCD163 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human sCD163 protein	Recombinant Human sCD163 protein
Information	Swissprot	Q86VB7	
Product details	Reactivity	Human	Human
	Host	Mouse	Mouse
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Protein A or G
	Specificity	Detects Human sCD163 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Human sCD163 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Human sCD163 Capture Antibody	
Capture			
ELISA Detection	1:1000-1:10000	Human sCD163 Detection Antibody (Biotin)	Optical Density
			0.01 0.1 0.1 0.1 0.1 0.1 0.1 1 0 100 10

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages, and may thereby protect tissues from free hemoglobin-mediated oxidative damage. This protein may also function as an innate immune sensor for bacteria and inducer of local inflammation. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.