A Reliable Research Partner in Life Science and Medicine

GAPDH Monoclonal Antibody

catalog number: E-AB-48016

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;Mouse;Rat;Bovine;Caprine;Guinea pig;Rabbit;Porcine; **Immunogen** Recombinant human GAPDH protein expressed by E.coli

 Host
 Mouse

 Isotype
 IgG1

 Clone
 3B3


Purification Protein A/G Purification

Buffer PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4

Applications Recommended Dilution

WB 1:5000-1:10000

Data




70— 44— 33— 26— 22—

Western Blot analysis of GAPDH in various cell lines at dilution of 1:5000. Lane1: DU145 whole cell lysate, Lane2: OS-RC-2 whole cell lysate, Lane3:T-47D whole cell Lysates, Lane4: HEC-1B whole cell lysate, Lane5: HepG2 whole cell lysate, Lane6: HEK-293 whole cell lysate, Lysates/proteins at 20g per lane.

Observed-MW:36 kDa Calculated-MW:36 kDa Western blot analysis of GAPDH in various tissues of different species at dilution of 1:5000. Lane1: Rat heart, whole tissue lysate, Lane2: Porcine brain, whole tissue lysate, Lane3: Bovine kidney, whole tissue lysate, Lane4:

Guinea pig liver, whole tissue lysate, Lane5: Rabbit pancreas, whole tissue ysate, Lane6: Caprine spleen, whole tissue lysate. Lysates/proteins at 20g per lane.

Observed-MW:36 kDa Calculated-MW:36 kDa



Immunofluorescent analysis of 293F cells using anti-GAPDH monoclonal antibody at 1:400 dilution.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com



Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

GAPDH is an enzyme of 37kDa that catalyzes the sixth step of glycolysis and thus serves to break down glucose for energy and carbon molecules. Because the GAPDH gene is often stably and constitutively expressed at high levels in most tissues and cells, it is commonly used as a loading control for western blot and as a control for qPCR.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V2.3