

A Reliable Research Partner in Life Science and Medicine

# PE/Cyanine7 Anti-Human CD74 Antibody[LL1]

Catalog Number: E-AB-F1421H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. LL1

Isotype Control PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792H]

Conjugation PE/Cyanine 7

Conjugation Information PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm

(e.g., a 780/60 nm bandpass filter).

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD19 Antibody and PE/Cyanine7 Anti-Human CD74 Antibody[LL1] (left) or PE/Cyanine7 Mouse IgG1,  $\kappa$  Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

### **Antigen Information**

 Uniprot ID
 P04233

 Gene ID
 972

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Background**

CD74 is a type II transmembrane glycoprotein also known as MHC class II associated invariant chain, invariant chain, Ii, MHC class II chaperone, and MIF receptor. CD74 exists in four isoforms with molecular masses of 33, 35, 41, and 43 kD, depending on genetic splicing. CD74 is primarily expressed on antigen presenting cells, including B cells, monocytes/macrophages, dendritic cells, and Langerhans cells. It is also expressed by activated T cells and activated endothelial and epithelial cells as well as carcinomas of lung, renal, gastric and thymic origin. The primary function of CD74 is intracellular sorting of MHC class II molecules and regulation of exogenous peptide loading onto MHC class II. It is also involved in the modulation of B cell differentiation and positive selection of CD4+ T cells. It has been reported that CD74 binds MIF (macrophage migration inhibitory factor) and signals through CD44 to regulate innate and adaptive immunity. It is also reported that H. pylori infection occurs through urease B binding of CD74 on gastric epithelial cells, inducing gastric epithelial cell apoptosis, NF-κB activation, and IL-8 production.

Fax: 1-832-243-6017