A Reliable Research Partner in Life Science and Medicine

Recombinant Human LRG1 Protein(His Tag)

Catalog Number: PDMH100450

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip			
	00	CPI	m	TO	m
ய	\mathbf{c}	UII	L L	ш	

Species Human

Source Mammalian-derived Human LRG1 protein Met1-Gln347, with an C-terminal His

 Mol_Mass
 38.0 kDa

 Accession
 P02750

Bio-activity Not validated for activity

Properties

Purity > 85% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis

Data



SDS-PAGE analysis of Human LRG1 proteins, 2µg/lane of Recombinant Human LRG1 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 43 kDa

Background

LRG1 (Leucine-Rich Alpha-2-Glycoprotein 1) is a Protein Coding gene. LRG1 belongs to the leucine-rich repeat (LRR) family. Members of this family are involved in protein-protein interaction, signal transduction, and cell adhesion and development. LRG1 is expressed during granulocyte differentiation. It contains 4 LIM zinc-binding domains and 1 Rho-GAP domain. LRG1 is involved in promoting neovascularization (new blood vessel growth) by causing a switch in transforming growth factor-beta (TGFbeta) signaling in endothelial cells. LRG1 binds to the accessory receptor endoglin and promotes signaling via the ALK1-Smad1/5/8 pathway. It may be a potential therapeutic target for the treatment of diseases where there is aberrant neovascularization. Diseases associated with LRG1 include Appendicitis and Normal Pressure Hydrocephalus.

For Research Use Only

Fax: 1-832-243-6017